

## **HEMATOPATHOLOGY REQUISITION**

Phone: (888) 261-2671 203 Indigo Dr., Brunswick, GA 31525 www.sepalabs.com



<u>Patient Information</u> (Please submit additional demographics separately.)	Client Information:
Name:	
DOB:	Date Performed:
	Biopsy performed by:
☐ Male ☐ Female	
Specimen Type	Send additional copy of report to:
☐ Bone Marrow ☐ Peripheral Blood ☐ Other:	Patient History (Please submit CBC and the last clinic note separately if available)
Materials Submitted	
Peripheral Smear #	
Aspirate Smear #	
Green Top Tube #	
Purple Top Tube #	
Culture Specimen:	
Other:	
DIAGNOSIS/DIFFERENTIAL DIAGNOSIS	
☐ Aplastic Anemia ☐ Lymphoma, Hodgkin	☐ B12 Deficiency
	gkin (specify if applicable):
☐ AML ☐ Myelodyslplastic Synd	drome ☐ Iron Deficiency
☐ CLL ☐ Myeloproliferative Ne	eoplasm (specify if applicable):
☐ CML ☐ Plasma Cell Dyscrasia	☐ Infectious Disease
☐ Other:	
SPECIAL STUDIES	
☐ COMPREHENSIVE ANALYSIS: Perform testing as necessary for diagnosis	
☐ Flow Cytometry	☐ PCR (Perform as necessary for diagnosis.)
☐ Fluid:	☐ BCR/ABL, Quantitative
☐ Leukemia/Lymphoma	☐ JAK-2 V617F
$\square$ PNH (peripheral blood specimen)	☐ JAK2 V617F-Reflex JAK2 Exon 12-14
☐ Tissue:	☐ JAK2 Exon 12-14
	☐ B cell Clonality
☐ Cytogenetics	☐ T cell Clonality
☐ FISH (Perform as necessary for diagnosis.)	☐ MPL
☐ CLL Panel	☐ Calreticulin
☐ MDS Panel	☐ Other:
☐ Prognostic Myeloma Panel	
☐ BCR/ABL	
☐ Other:	
SPECIMEN REQUIREMENTS	
Bone Marrow: For comprehensive bone marrow biopsy evaluation, please submit at least 2 mL of aspirate in EDTA/purple top and at least 2 mL of aspirate in	
sodium heparin/green top. If no aspirate is obtained (dry tap), core biopsy may be submitted in RPMI for ancillary testing (in addition to the core biopsy submitted in formalin for histologic evaluation)	

PCR: 2-5 mL of peripheral blood (or 1-2 mL bone marrow aspirate) In EDTA/purple top. Refrigerate/transport with ice pack. Do not freeze.

FISH: 2-5 mL of peripheral blood (or 1-2 mL bone marrow aspirate) in sodium heparin/green top (preferred) or EDTA/purple top or tissue in RPMI. Refrigerate/transport with ice pack. Do not freeze.

Flow Cytometry: At least 3 mL of peripheral blood (or 1-2 mL bone marrow aspirate) in EDTA/purple top or sodium heparin/green top. Refrigerate/transport with ice pack. Do not freeze.